

Hyperon–Nucleon Final State Interaction in Kaon Photoproduction of the Deuteron

B.O.Kerbikov

Institute of Theoretical and Experimental Physics,
117218, Moscow, Russia

Abstract

Final state hyperon-nucleon interaction in strangeness photoproduction of the deuteron is investigated making use of the covariant reaction formalism and the P -matrix approach to the YN system. Remarkably simple analytical expression for the amplitude is obtained. Pronounced effects due to final state interaction are predicted including the manifestation of the 2.13 GeV resonance.

1 Introduction

Up to now most of the information on the hyperon–nucleon (YN) interaction has been obtained either from hypernuclei or from K^-d and π^+d reactions. After several decades of studies our knowledge on the YN system is still far from being complete.

Recently the interest to the $\Lambda N - \Sigma N$ system has flared again in connection to the expected CEBAF experimental results on the kaon photoproduction on the deuterium [1]. The final state YN interaction (FSI) plays an important role in the $\gamma d \rightarrow K^+ Y n$, $Y = \Lambda, \Sigma^0$ reaction. Therefore high resolution photoproduction experiments can substantially deepen our understanding of the YN dynamics.

The problem of FSI in $\gamma d \rightarrow K^+ Y n$ reaction has been addressed by several authors starting from the pioneering paper by Renard and Renard [2,3,4]. Two novel features differ the present work from the previous studies. First, covariant formalism based on direct evaluation of Feynman diagrams is used which allows to analyse the data beyond the region of low energy and low momentum transfer. Second, the YN interaction is described within the P -matrix approach which takes into account the subnuclear degrees of freedom and disentangle the dynamical singularities from kinematical threshold effects [5]. The P -matrix analysis of the YN interaction was presented in [6] (see also [7]) and we shall use the set of parameters from [6].

In our previous publications [8],[9] we have announced some preliminary results without proving the central assertion, namely that the FSI effects allow remarkably simple evaluation within the P -matrix approach. In the present paper we give a detailed derivation of this result. Our central result is that YN FSI in strangeness photoproduction of the deuteron is quite significant and that P -matrix provides a remarkably transparent approach to this phenomenon.

The organization of the paper is the following. In Section 2 we present kinematical relations relevant for the reaction and display equations for the cross sections. Analysis of the plane waves approximation is given in Section 3. In Section 4 we present a summary of the P -matrix approach to the hyperon-nucleon FSI. In Section 5 the loop diagram with YN FSI is calculated and expressed through the enhancement factor in the spirit of the Migdal–Watson FSI theory. Numerical results and calculations are

presented in Section 6.

2 Kinematics and Cross-section

The reaction $\gamma d \rightarrow K^+ Y n$, $Y = \Lambda, \Sigma^0$ is a $2 \rightarrow 3$ process. The corresponding kinematical relations as well as expressions for various cross-sections and distributions are well known (see e.g. [10]). However we find it appropriate to include a list of basic equations in order to facilitate comparison with other theoretical papers and to provide a framework for the presentation of the experimental results.

We start with a general expression for the $2 \rightarrow 3$ reaction

$$d\sigma = (2\pi)^4 \delta^{(4)}(k + p_d - p_K - p_Y - p_n) \frac{|T|^2}{4I} \frac{d^3 p_K}{(2\pi)^9 2E_K} \frac{d^3 p_Y}{2E_Y} \frac{d^3 p_n}{2E_n}, \quad (2.1)$$

where k is the four-momentum of the initial photon, I is the flux factor which in the deuteron rest frame is equal to kM_d , and the Dirac spinors are normalized according to $\bar{u}u = 2m$. The number of essential final state Lorentz scalar variables for 3-body final state is 4 with the standard set being

$$s_1 = (p_K + p_Y)^2, s_2 = (p_Y + p_n)^2, t_1 = (k - p_K)^2, t_2 = (p_d - p_n)^2. \quad (2.2)$$

In addition the total energy squared $s = (k + p_d)^2 = (p_K + p_Y + p_n)^2$ will also be used in equations to follow.

The physical quantity we are interested in is the double differential cross-section

$$d^2\sigma \equiv \frac{d^2\sigma}{d|\mathbf{p}_K|d\Omega_K} = \frac{1}{2^{11}\pi^5 k M_d E_K} \frac{\lambda^{1/2}(s_2, m_Y^2, m_n^2)}{s_2} \int d\Omega_{Yn}^* |T|^2. \quad (2.3)$$

Here k , $\mathbf{p}_K^2$, E_K and Ω_K correspond to the deuteron rest system (LS) with z -axis defined by the incident photon beam direction $\mathbf{k}$. The solid angle Ω_{Yn}^* is defined in the Yn center-of-momentum system (CMYn). The quality $\lambda(x, y, z)$ is the standard kinematical function $\lambda(x, y, z) = x^2 - 2(y+z)x + (y-z)^2$. Integration over $d\Omega_{Yn}^*$ can be replaced by integration over $dt_2 ds_1$ since linear relations hold between t_2 and $\cos\theta^*$ and s_1 and $\cos\varphi^*$ correspondingly. With such a replacement we can write $d^2\sigma$ in terms of the Chew-Low plot as

$$\frac{d^2\sigma}{ds_2 dt_1} = \frac{1}{2^{10}\pi^4 \lambda(s, 0, M_d^2)} \int dt_2 ds_1 \frac{|T(t_2, s_1)|^2}{[-\Delta_4(t_1, s_1, s_2, t_2)]^{1/2}}, \quad (2.4)$$

where Δ_4 is a 4×4 symmetric Gram determinat (see [10] for an explicit expression) which is a quadratic in any of the variables t_1, s_1, s_2, t_2 . The region of integration in equation (2.4) has to satisfy $\Delta_4 \leq 0$. For s_2 and t_1 fixed (which corresponds to fixed $|\mathbf{p}_k|$ and Ω_K) the boundary of integration $\Delta_4 = 0$ can be solved for s_1 in terms of t_2 :

$$s_1^\pm = m_K^2 + m_Y^2 - \frac{1}{2t_1} \{ (t_1 + m_K^2)(t_1 + m_Y^2 - t_2) \pm \lambda^{1/2}(t_1, m_K^2, 0) \lambda^{1/2}(t_1, m_Y^2, t_2) \}. \quad (2.5)$$

Expressed in terms of $s_1^\pm$, the function Δ_4 reads

$$16\Delta_4 = \lambda(s_2, t_1, M_d^2)(s_1 - s_1^+)(s_1 - s_1^-), \quad (2.6)$$

Actual calculations presented below were performed making use of (2.4) since integration over invariant variables t_2 and s_1 is more straightforward than integration over the angle Ω_{Yn}^* needed in (2.3).

3 The Plane Waves Approximation

We shall use covariant relativistic approach to calculate the amplitude T of the $\gamma d \rightarrow K^+ Y n$ process. The amplitude will be approximated by the contributions of the two diagrams, namely the tree (pole, or plane waves) graph and the triangle graph with Yn FSI. It will be shown that within approximations to be specified below the sum of these diagrams reduces to the plane waves diagram multiplied by an enhancement factor depending on s_2 and therefore factorizing out of the integral in (2.4)

To calculate the plane-waves diagram two blocks have to be specified: (i) the elementary photoproduction amplitude $M^{\gamma K}$ on the proton, and (ii) the deuteron vertex Γ_d . The elementary amplitude used in the present calculation was derived from the tree level effective Lagrangian with the account of several resonances in s, t and u channels [11]. This amplitude has the following decomposition over invariant terms [12]

$$M^{\gamma K} = \bar{u}_Y \sum_{j=1}^6 \mathcal{A}_j \mathcal{M}_j(s', t', u') u_p, \quad (3.1)$$

where

$$s' = (k + p_p)^2, \quad t' = (k - p_K)^2, \quad u' = (k - p_Y)^2. \quad (3.2)$$

The decomposition of the deuteron vertex function Γ_d in independent Lorentz structures has the form [13]

$$\Gamma_d = \sqrt{m_N} [(p_p + p_n)^2 - M_d^2] \times \left[\varphi_1(t_2) \frac{(p_p - p_n)_\mu}{2m_N^2} + \varphi_2(t_2) \frac{1}{m_N} \gamma_\mu \right] \mathcal{E}^\mu(p_d, \lambda). \quad (3.3)$$

Here $\mathcal{E}^\mu(p_d, \lambda)$ is the polarization 4-vector of the deuteron with momentum p_d and polarization λ . The vertex (3.3) implies that the deuteron as well as the spectator neutron are on mass shell while the proton is off its mass shell. In the nonrelativistic limit (but with relativistic corrections) the functions φ_1 and φ_2 may be expressed in terms of the deuteron S - and D -waves [14]

$$\varphi_1 = \frac{m_N^2}{2\mathbf{p}_n^2} \left[-\left(1 + \frac{m_N}{2E_n}\right) u_D + \frac{1}{\sqrt{2}} \left(1 - \frac{m_N}{E_n}\right) u_S \right], \quad (3.4)$$

$$\varphi_2 = \frac{m_N}{2E_n} (\sqrt{2} u_S + u_D), \quad (3.5)$$

with $E_n = \sqrt{\mathbf{p}_n^2 + m_N^2}$. The normalization condition is the following

$$\frac{m_N}{2\pi^2} \int_0^\infty \frac{dp_n \mathbf{p}_n^2}{E_n} [u_S^2(p_n) + u_D^2(p_n)] = 1. \quad (3.6)$$

Now we can write the following expression for the tree diagram

$$T^{(t)} = \bar{u}_Y \left\{ \left(\sum_{j=1}^6 \mathcal{A}_j \mathcal{M}_j(s', t', u') \right) S(p_p) \Gamma_d \right\} u_n^c, \quad (3.7)$$

where $S(p_p)$ is the proton propagator and u_n^c is a charge conjugated neutron spinor. The deuteron vertex in (3.7) may be substituted by the relativistic deuteron wave function according to [15]

$$\psi_d = [2(2\pi)^3 M_d]^{1/2} S(p_p) \Gamma_d. \quad (3.8)$$

Then (3.7) can be rewritten in the following schematic form

$$T^{(t)} = [2(2\pi)^3 M_d]^{1/2} M^{\gamma K} \psi_d, \quad (3.9)$$

where ψ_d is the relativistic deuteron wave function discussed at length in [15], and where summation over magnetic quantum numbers is tacitly assumed.

In the numerical calculations of the plane waves diagram use has been made of the Gross relativistic deuteron wave function [15]. Finally we

mention that nonrelativistic reduction of Eqs. (2.3), (2.4) and (3.9) is straightforward and yields the standard impulse approximation

$$\frac{d^2\sigma}{d|\mathbf{p}_K|d\Omega_K} = \frac{\mathbf{p}_K^2 |\mathbf{p}_Y^*|}{64\pi^2 k E_K \sqrt{s_2}} \int d\Omega_{Yn}^* |M^{\gamma K}|^2 |\psi_d^{nr}|^2, \quad (3.10)$$

where the nonrelativistic deuteron wave function normalization condition is

$$\int d^3p_n |\psi_d^{nr}(p_n)|^2 = 1. \quad (3.11)$$

4 Hyperon-Nucleon interaction in the P -matrix approach

Before we consider the triangle diagram with Yn rescattering it is worth to recapitulate the main results of the P -matrix approach to hyperon-nucleon interaction [6]. Since FSI is essential only at low relative momenta of the Yn pair the nonrelativistic treatment is quite legitimate.

First we recall the interpretation of the P -matrix in terms of the underlying coupled-channel quark-hadron potential [16]. Namely, the Jaffe-Low P -matrix description is equivalent to the coupling between hadron and quark channels via the nonlocal energy dependent potential of the form

$$V_{\text{hqh}} = \sum_n \frac{f_n(r) f_n(r')}{E - E_n}, \quad (4.1)$$

where E_n are the energies of the six-quark "primitives" [17,5] (P -matrix poles), and the form factors $f_n(r)$ are given by

$$f_n(r) = \frac{c_n}{b\sqrt{4\pi}} \delta(r - b). \quad (4.2)$$

Here b is the bag radius [17,16,5] and $|c_n|^2 = \lambda_n^2 / 2\mu_{Yn}$, where λ_n are the residues of the P -matrix [17,16,5]

$$P = P_0 + \sum_n \frac{\lambda_n^2}{E - E_n}. \quad (4.3)$$

As it was shown in [6] a single primitive at $E_n = 2.34$ GeV is sufficient for a high quality description of the existing YN experimental data, so that the subscript n will be omitted. To avoid lengthy expressions (4.3) is written for the case of a single hadron channel, namely ΛN , inclusion of

the $\Lambda N - \Sigma N$ conversion is straightforward [6] and the final results with the account of the ΣN channel will be presented below.

The scattering problem corresponding to the interaction (4.1),(4.2) can be easily solved. In particular for the momentum space half-of-shell amplitude $F(p, p'; E')$, $E' = p'^2/2\mu_{Yn} \neq p^2/2\mu_{Yn}$ and for the Jost function $f(-p')$ one gets

$$F(p, p'; E') = -\frac{2\pi}{\mu} \lambda^2 b^2 \frac{\sin pb}{pb} d^{-1}(E') \frac{\sin p'b}{p'b}, \quad (4.4)$$

$$d(E') = E' - E_n + \frac{\lambda^2}{p'} e^{ip'b} \sin p'b, \quad (4.5)$$

$$f^{-1}(-p') = \frac{E' - E_n}{E' - E_n + \frac{\lambda^2}{p'} e^{ip'b} \sin p'b}. \quad (4.6)$$

These results will be used in the next section devoted to the calculation of the triangle diagram with Yn FSI.

5 Triangle Diagram

Next we consider the triangle (loop) diagram with Yn FSI. The general expression for this graph reads

$$T^{(l)} = \int \frac{d^4 p_n}{(2\pi)^4} \bar{u}_Y(p'_Y) \left\{ \left(\sum_{j=1}^6 \mathcal{A}_j \mathcal{M}_j \right) S(p_p) \Gamma_d C S(p_n) T_{Yn} S(p_Y) \right\} \bar{u}_n(p'_n). \quad (5.1)$$

Here C is the charge conjugation matrix, T_{YN} is a hyperon-nucleon vertex, this vertex being “dressed” by the corresponding spinors constitutes the hyperon-nucleon amplitude F (see (4.4)).

The comprehensive treatment of the triangle diagram is a purpose of the forthcoming publication while here we resort to some approximations with the aim to expose the effects of the FSI. Only positive frequency components are kept in the propagators $S(p_n)$ and $S(p_Y)$ in (5.1), while the propagator $S(p_p)$ together with Γ_d is lumped into the relativistic deuteron wave function ψ_d [15]. Then integration over the time component dp_n^0 is performed. Thus we arrive at the following expression for $T^{(l)}$

$$T^{(l)} = 2\mu[(2\pi)^3 2M_d]^{1/2} \int \frac{d\mathbf{p}_n}{(2\pi)^3} \frac{M^{\gamma K} \psi_d(p_n) F(p, p'; E')}{\mathbf{p}^2 - \mathbf{p}'^2 - i0}, \quad (5.2)$$

where $\mathbf{p}$ and $\mathbf{p}'$ are the YN c.m. momenta before and after the FSI, $\mathbf{p} = \mathbf{p}_n - \frac{1}{2}(\mathbf{k} - \mathbf{p}_K)$. The quantity $F(p, p'; E')$ is the half-off-shell Yn amplitude given by (4.4)-(4.5). The arguments of the elementary photoproduction amplitude $M^{\gamma K}$ in (5.2) are specified by (3.1)-(3.2) but in the kinematical region where FSI is essential $M^{\gamma K}$ can be considered as point-like and hence factored out of the integral (5.2) at the values of the arguments fixed by the energies and momenta of the initial and final particles (i.e. at the values of the arguments corresponding to the plane-wave diagram). Thus (5.2) takes the form

$$T^{(l)} = -4\pi\lambda^2 b^2 [(2\pi)^3 2M_d]^{1/2} d^{-1}(E') \frac{\sin p'b}{p'b} M^{\gamma K} J(p'), \quad (5.3)$$

where

$$J(p') = \int \frac{d\mathbf{p}_n}{(2\pi)^3} \frac{\sin pb}{pb} \frac{\psi_d(p_n)}{\mathbf{p}^2 - \mathbf{p}'^2 - i0}. \quad (5.4)$$

In the complex p plane integral (5.4) picks up contributions from the poles of the propagator and from the singularities of the deuteron wave function. The dominant contribution stems from the poles of the propagator. Numerical calculations using realistic deuteron wave function will be presented elsewhere. Here we present simple arguments illustrating the above conclusion. Consider zero-range deuteron wave function which in the momentum space has the form

$$\psi_d(p_n) = \frac{\sqrt{\alpha}}{\pi(\mathbf{p}_n^2 + \alpha^2)}, \quad (5.5)$$

where $\alpha = \sqrt{m_n|E_B|} \simeq 45.7$ MeV. Recalling that $\mathbf{p}_n = \mathbf{p} + \mathbf{Q}/2$, $\mathbf{Q} = \mathbf{k} - \mathbf{p}_K$, and performing in (5.4) the angular integration, one gets

$$\begin{aligned} J(p') &= \frac{\sqrt{\alpha}}{\pi} \int_0^\infty \frac{dp p^2}{(2\pi)^3} \frac{\sin pb}{pb} (\mathbf{p}^2 - \mathbf{p}'^2 - i0)^{-1} \frac{2\pi}{pQ} \ln \left| \frac{(p + \frac{Q}{2})^2 + \alpha^2}{(p - \frac{Q}{2})^2 + \alpha^2} \right| \simeq \\ &\simeq \frac{16\sqrt{\alpha}}{Q} \int_0^\infty \frac{dp p^2}{(2\pi)^3} \frac{\sin pb}{pb} (\mathbf{p}^2 - \mathbf{p}'^2 - i0)^{-1} (1 - \frac{4\alpha^2}{Q^2}). \end{aligned} \quad (5.6)$$

In making the approximation (5.6) use was made of the fact that FSI is effective at $pb \lesssim 1$ while the typical momentum transfer is $Q \gg 1/b$ and $Q \gg \alpha$. Formally this is reflected in the fact that the integral $J(p')$ is damped at high values of p due to the formfactor $\sin pb/pb$. the remaining

integral (5.5) is trivial and yields

$$J(p') = \left(\frac{4\sqrt{\alpha}}{\pi Q^2}\right) \frac{e^{ip'b}}{4\pi b} \left(1 - \frac{4\alpha^2}{Q^2}\right) \simeq \psi_d(p_n) \frac{e^{ip'b}}{4\pi b} \left(1 - \frac{4\alpha^2}{Q^2}\right). \quad (5.7)$$

Then using (5.3) and (4.6) one gets for the loop diagram

$$T^{(l)} = T^{(t)} \left[-1 + \frac{1}{f(-p')} \left(1 - \frac{4\alpha^2 c}{Q^2}\right)\right]. \quad (5.8)$$

Here $T^{(t)}$ is the plane-waves amplitude (3.9), $f(-p')$ is the Jost function given by (4.6) and

$$c = \frac{\lambda^2 b}{E_n - E} \frac{\sin p'b}{p'b}.$$

Within few percent accuracy the term $4\alpha^2 c^2/Q^2$ can be neglected since $Q^2 \gg \alpha^2$ and $c \ll 1$. The last inequality is based on the set of parameters fitted to the YN interaction [6], namely $\lambda^2 = 0.014 \text{ GeV}^2$ and $E_n = 2.34 \text{ GeV}$. For this set of parameters $c \lesssim 0.2$ in the region where FSI is important.

Therefore for the sum of the plane waves and loop diagrams one gets

$$T^{(t)} + T^{(l)} \simeq T^{(t)} / f(-p') \quad (5.9)$$

Thus we arrived at a remarkably simple and physically transparent result in the spirit of the Migdal-Watson FSI theory. This result might seem trivial since it can be immediately obtained by factoring out the deuteron wave function¹ out of the integral (5.4). However apriori such an approximation does not look reliable and has to be justified making us of the amplitude corresponding to the P -matrix as it was done above. Analogous derivation with the realistic deuteron wave function will be presented in the forthcoming publication. Inclusion of the $\Lambda N - \Sigma N$ conversion in the P -matrix approach is very simple [6]. Namely, in (5.9) the two-channel Jost function has to be used. It is given by

$$F^{-1}(-p'_1, -p'_2) = \frac{E' - E_n}{E' - E_n + \frac{\lambda_1^2}{P_1^0 - ip'_1} + \frac{\lambda_2^2}{P_2^0 - ip'_2}}, \quad (5.10)$$

where the indices 1,2 correspond to ΛN and ΣN channels and the values of the parameters were fitted in [6]. Then the final expression for the

¹At the value of p_n corresponding to the tree diagram with the same kinematical variables of the initial and final particles.

amplitude corresponding to the sum of the plane waves and loop diagrams with couple-channel FSI included reads

$$T^{(t)} + T^{(l)} \simeq T^{(t)} / F(-p'_1, -p'_2). \quad (5.11)$$

Thus using simple arguments we have splitted the description of the $\gamma d \rightarrow K^+ Y n$ reaction into two blocks: (i) the covariant calculation of the plane waves diagram with relativistic deuteron wave function and relativistic elementary photoproduction amplitude, and (ii) the calculation of the P -matrix enhancement factor with a set of parameters fitted to the existing data on YN interaction. The accuracy of this factorization is of the order of few percent which can be seen from model considerations presented above and will be confirmed by numerical calculations in the forthcoming publication.

6 Results and Conclusions

Here we present the results of the calculations performed according to Eqs. (2.4), (3.9), (5.10) and (5.11). In Fig. 1 we plot the double differential cross-section (2.3) at $|\mathbf{p}_k| = 1.091$ GeV and $\Theta_K = 0^\circ$ in a rather wide range of the photon energy covering the ΣN threshold region with nearby 2.13 GeV resonance. The behavior of the cross section is very instructive. Consider first the dashed line corresponding to the plane waves approximation. At this particular kinematics ($P_K = 1.091$ GeV, $\Theta_{\gamma K} = 0^\circ$) the threshold of Λn production is at $E_\gamma = 1.4$ GeV and according to (3.10) the factors that governs the near threshold cross section behavior are the phase space and the momentum dependence of the deuteron wave function. The sigma-nucleon threshold in Fig.1 corresponds to $E_\gamma = 1.5$ GeV. The ΣN cusp in the cross section is practically invisible. This fact was known long ago – see e.g. [18]. It is a typical situation since "In the experimental cross sections of nuclear reactions the cusps are rarely seen" [19]. To see a cusp one should accurately take into account FSI and moreover in order to be pronounced the cusp should be enhanced by nearby pole like 2.13 GeV resonance close to ΣN threshold. Next we turn to the solid line in Fig. 1 with FSI included. The ΛN FSI interaction causes the cross section to rise more rapidly close to threshold as was noticed before in [3]. Then one observes a very clear signal at the ΣN threshold ($E_\gamma = 1.49$) GeV caused by the pole at 2.13 GeV. We remind that according to the analysis

of [6,7] the 2.13 GeV resonance is not a genuine six-quark state but a P -matrix partner of the deuteron. Our prediction that this resonance can be investigated in strangeness photoproduction reaction is at odds with the conclusions made in [3] that FSI is important only close to Λn threshold and is completely insignificant at higher energies. The attribution of the pole corresponding to the 2.13 GeV resonance to a particular sheet of the multi-sheeted Riemann surface is a long-standing problem [6] and our hope is that photoproduction experiments can shed a new light on it. Concerning the structure close to ΣN threshold another remark has to be made. Namely, one may ask whether such structure can be generated simply by the triangle graph with constant $\Lambda N - \Sigma N$ conversion matrix element. This question has been addressed in the literature for the case of a similar $K^- d \rightarrow \pi^- \Lambda p$ reaction [20,21], and it was shown that the singularity of the triangle diagram by itself can not explain the observed structure. Finally a remark is due on another approximation tacitly made in the present work. This concerns hadronic corrections like pion-exchange to the basic P -matrix YN interaction as well as corrections due to nonorthogonality of the six-quark and YN states [16]. As was shown in [6] the present experimental data on YN interaction are not sufficient to feel these corrections. From the theoretical side such corrections modify the Jost function given by (4.6) and (5.11), in particular they smear the zero of the inverse Jost function at $E = E_n$.

The author would like to thank V.A.Karmanov for many fruitful discussions and suggestions. Help in numerical calculations from N.O.Agasian is gratefully acknowledged. This work was possible due to stimulating contacts with C.Fayard, G.H.Lamot, F.Rouvier and B.Saghai. Hospitality and financial support from the University Claude Bernard and DAPNIA (Saclay) are gratefully acknowledged.

References

- [1] B. Mecking et al., CEBAF Proposal PR-89-045, (1989).
- [2] F.M.Renard and Y.Renard, Nucl. Phys. **B1**, 389 (1967).
- [3] R.A.Adelseck, L.E.Wright, Phys. Rev. **C39**, 580 (1989).
- [4] X.Li, L.E.Wright, J.Phys. **G17**, 1127 (1991);
S.R.Cotanch, S.Hsiao, Nucl. Phys. **A450**, 419c (1986);
T.-S.H.Lee et al., Nucl. Phys. **A639**, 247c (1998);
H.Yamamura et al., Nucl.-th/9907029 V.2.
- [5] S.V.Bashinsky, R.L.Jaffe, Nucl. Phys. **A625**, 167 (1997).
- [6] B.L.G.Bakker et al., Sov. J. Nucl. Phys. **43**, 982 (1986);
B.Kerbikov et al., Nucl. Phys. **A480**, 585 (1988).
- [7] L.Kisslinger and D.P.Heddle, Phys. Rev. **C30**, 965 (1984).
- [8] B.Kerbikov, hep-ph/9910361.
- [9] B.Kerbikov, hep-ph/003054.
- [10] E.Byckling and K.Kajantie, Particle Kinematics (John Willey and Sons, London-N.Y., 1973).
- [11] J.C. David J.C. et al., Phys. Rev. **C53**, 2613 (1996).
- [12] G.F.Chew et al., Phys. Rev. **106**, 1345 (1957).
- [13] M.Gourdin et al., Nuovo Cim. **37**, 525 (1965).
- [14] V.A.Karmanov and A.V.Smirnov, Nucl. Phys. **A575**, 520 (1994).
- [15] F.Gross, Phys. Rev. **D10**, 223 (1974);
W.W.Buck, F.Gross, Phys. Rev. **D20**, 2361 (1979).
- [16] Yu.A.Simonov, Phys. Lett. **B1071**, 1 (1981);
Yu.A.Simonov, Nucl. Phys. **A463**, 231c (1987).
- [17] R.L.Jaffe, F.E.Low, Phys. Rev. **D19**, 2105 (1979).
- [18] F.H.Renard and Y.Renard, Phys. Lett. **24B**, 159 (1967).

- [19] A.M.Badalyan, L.P.Kok, M.I.Polikarpov, Yu.A.Simonov, Phys. Rep. **82** , 31 (1982).
- [20] H.G.Dosch and I.O.Stamatescu, Z.Phys. **C3**, 249 (1980).
- [21] C.Pigot et al., Nucl. Phys. **B249**, 172 (1985).

Figure Caption

Fig.1 The double differential cross section as a function of the photon energy for $p_K = 1.091$ GeV, $\Theta_{\gamma K} = 0^\circ$. Dashed line corresponds to the plane waves approximation, solid line – with Yn FSI included.